


# Compare Pairs: Bodyfat Before, After

Data A1:B32

Last updated 2 February 2016 at 12:21 by Analyse-it Software, Ltd.

## Descriptives




Bodyfat	N	Mean	Mean SE	SD
Before	28	23.81	1.607	8.50
After	28	19.14	1.175	6.22
Differences	28	4.68	0.547	2.89

Data A1:B32

Last updated 2 February 2016 at 12:21 by Analyse-it Software, Ltd.

Differences



Location

Hodges-Lehmann shift | -4.75  
 95.49% CI | -5.90 to -3.55

$$F(\text{Bodyfat: After}) = F(\text{Bodyfat: Before} + \Delta)$$

Wilcoxon test

Hypothesized difference | 0

Sign	n	Rank sum	Mean rank
Positive	2	4.5	2.25
Negative	26	401.5	15.44
Zero	0		

T statistic | 4.50  
 Z approximation | -4.52  
 p-value | <0.0001<sup>1</sup>

H0:  $\Delta = 0$

The shift in location between the distributions of the populations is equal to 0.

H1:  $\Delta \neq 0$

The shift in location between the distributions of the populations is not equal to 0.

<sup>1</sup> Reject the null hypothesis in favour of the alternative hypothesis at the 5% significance level.